

β-肌动蛋白/β-Actin (内参) 抗体

<p>Host: Rabbit</p> <p>Target Protein: beta-Actin (Loading Control)</p> <p>IR: Immunogen Range:1-200/375</p> <p>Clonality: Polyclonal</p> <p>Isotype: IgG</p> <p>Entrez Gene: 60</p> <p>Swiss Prot: P60709</p> <p>Source: Synthetic MAP peptide derived from human beta-Actin:1-200/375</p> <p>Purification: affinity purified by Protein A</p> <p>Storage: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Loading Control This gene encodes one of six different actin proteins. Actins are highly conserved proteins that are involved in cell motility, structure, and integrity. This actin is a major constituent of the contractile apparatus and one of the two nonmuscle cytoskeletal actins. [provided by RefSeq, Jul 2008].</p>	<p>Size: 50ul 100ul</p> <p>Concentration: 1mg/ml</p> <p>Applications: WB(1:5000-50000) ELISA(1:5000-20000) IHC-P(1:100-500) Flow-Cyt(1µg/Test) ICC(1:100)</p> <p>Cross Reactive Species: Human Mouse Rat Chicken Dog Pig Rabbit Sheep Bee Fish Guinea Pig Hamster Cat 3 2 .</p> <p>For research use only. Not intended for diagnostic or therapeutic use.</p>
--	---