

## KIF18B Rabbit pAb

Catalog Number: BN40692R

Target Protein: KIF18B

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Rat (predicted:Human,Mouse,Sheep,Cow,Dog,Horse)

Predicted MW: 94 kDa

Entrez Gene: 146909

Swiss Prot: Q86Y91


Source: KLH conjugated synthetic peptide derived from human KIF18B : 121-220/864.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** The Kinesins constitute a large family of microtubule-dependent motor proteins which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Individual Kinesin members play crucial roles in cell division, intracellular transport and membrane trafficking events, including endocytosis and transcytosis. KIF18B (kinesin family member 18B) is an 864 amino acid nuclear and cytoplasmic protein that belongs to the kinesin-like protein family. Containing one kinesin-motor domain, KIF18B is dominantly expressed in amygdala. During mitosis, KIF18B is suggested to be closely associated with astral microtubules emanating from the spindle pole during prometaphase and metaphase. KIF18B may play an important role in regulating cell division. KIF18B exists as four alternatively spliced isoforms and is encoded by a gene located on human chromosome 17q21.31.



Paraformaldehyde-fixed, paraffin embedded (rat lymphoid); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KIF18B) Polyclonal Antibody, Unconjugated (BN40692R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) instructions and DAB staining.