


ZDHHC6 Rabbit pAb

BN40522R


DATASHEET

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 64429	SWISS: Q9H6R6	
Target: ZDHHC6		
Immunogen: KLH conjugated synthetic peptide derived from human ZDHHC6: 221-320/413.		Reactivity: Human, Mouse, Rat (predicted: Pig, Horse)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 48 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cell membrane


VALIDATION IMAGES




Sample: Kidney (Mouse) Lysate at 40 ug Primary: Anti-ZDHHC6 (BN40522R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kD
Observed band size: 48 kD



Sample: Lane 1: Mouse Pancreas tissue lysates Lane 2: Mouse Cerebrum tissue lysates Lane 3: Mouse Testis tissue lysates Lane 4: Mouse Raw264.7 cell lysates Lane 5: Mouse NIH/3T3 cell lysates Lane 6: Rat Kidney tissue lysates Lane 7: Rat Cerebrum tissue lysates Lane 8: Rat Testis tissue lysates Lane 9: Human HL-60 cell lysates
Primary: Anti-ZDHHC6 (BN40522R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kDa Observed band size: 50 kDa



Paraformaldehyde-fixed, paraffin embedded (mouse placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (ZDHHC6) Polyclonal Antibody, Unconjugated (BN40522R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (instructions and DAB staining).



Paraformaldehyde-fixed, paraffin embedded (rat placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (ZDHHC6) Polyclonal Antibody, Unconjugated (BN40522R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) instructions and DAB staining.

本产品仅用于科研

TEL: 010-82422342 www.biorigin.Ltd